

OVERVIEW OF APPLICATION REQUIREMENTS

Application requirements for XtendiMax® herbicide with VaporGrip® Technology, a restricted use pesticide, are intended to help maximize weed control with on-target applications and minimize the potential of off-target movement.

MANDATORY TRAINING & CERTIFICATION

Only applicators who are certified and have completed dicamba or auxin-specific training may apply.

APPLICATION TIMING

Applications must occur between one hour after sunrise and two hours before sunset. (DO NOT spray during an INVERSION.)

APPLICATOR PREPARATION

RECORD KEEPING

Certified applicator MUST fill out records within 72 hours of spraying and keep them for a period of 2 years.

SENSITIVE CROPS

Do not apply when wind is blowing toward adjacent sensitive crops.

ENDANGERED SPECIES CONCERNS

Additional protection measures are required in specific counties where endangered species may exist. Visit EPA.gov/Endangered-Species or call 1-844-447-3813 for more information.

DOWNWIND BUFFER

Maintain the required label buffer. (Minimum 110 ft buffer at 22 fl oz/A rate.)

SPRAYER SETUP

☐ SPRAY VOLUME

Apply in a minimum of 15 gallons of spray solution per acre.

□ GROUND SPEED

Do not exceed 15 mph ground speed.

☐ SPRAY BOOM HEIGHT

Do not exceed a boom height of 24 inches above target pest or crop canopy.

■ WIND SPEED

Apply when wind speed, measured at boom height, is between 3 and 10 mph. (DO NOT spray during an INVERSION.)

☐ SPRAY SYSTEM EQUIPMENT CLEANOUT

Ensure that entire sprayer system is properly cleaned before AND after using this product to avoid potential contamination.

NOZZLES & TANK MIXES

□ NOZZLES

Use only approved nozzles within specified pressure.

□ TANK-MIX PARTNERS

Use only approved tank-mix partners. *Ammonium sulfate* and ammonium-based additives are prohibited in applications that include XtendiMax with VaporGrip Technology.

Visit XtendiMaxApplicationRequirements.com for approved tank-mix partners, nozzles and label.

APPLICATION RATES*

COMBINED TOTAL PER YEAR FOR ALL APPLICATIONS	88 FLUID OUNCES PER ACRE
TOTAL OF ALL BURNDOWN/EARLY PREPLANT, PREPLANT, AT PLANTING, AND PRE-EMERGENCE APPLICATIONS	44 FLUID OUNCES PER ACRE
TOTAL OF ALL POSTEMERGENCE (IN-CROP) APPLICATIONS	44 FLUID OUNCES PER ACRE
MAXIMUM POSTEMERGENCE (IN-CROP), SINGLE APPLICATION	22 FLUID OUNCES PER ACRE

* These rates are for Roundup Ready 2 Xtend® soybeans and cotton with XtendFlex® Technology

XtendiMax® herbicide with VaporGrip® Technology Application Window in Soybeans (Apply prior to beginning bloom or no more than 45 days after planting, whichever comes first) Pre-emergence Beginning Bloom (R1) Pre-emergence VE VC V1 V2 V3 R1 R3 XtendiMax® herbicide with VaporGrip® Technology Application Window in Cotton (Apply up to mid-bloom stage or no more than 60 days after planting, whichever comes first) Pre-emergence Mid-Bloom Mid-Bloom Harvest

** For any other use of XtendiMax® herbicide with VaporGrip® Technology please refer to product label at XtendiMaxApplicationRequirements.com

THIS SUMMARY IS NOT A SUBSTITUTE FOR READING AND FOLLOWING ALL PRODUCT LABELING.

FOR PRODUCT QUESTIONS OR INQUIRIES AND/OR TO REPORT ANY NON-PERFORMANCE OF THIS PRODUCT AGAINST ANY PARTICULAR WEED SPECIES, CALL 1-844-RRXTEND